

UbiQ

targeting the ubiquitin system

Ub-Ub(20-33)-FP K27 linked *(human sequence, synthetic)*

UbiQ code : UbiQ-045
Batch # : B01012013-001
Amount : 25 ug, 1 uL of 2.38 mM stock in DMSO
Purity : ≥95% by RP-HPLC
Mol. Weight : 10.5 kDa
Storage : upon arrival, store solution at –80°C. Store dark and avoid multiple freeze/thaw cycles.

Productsheet

Background. UbiQ-045 is a fluorescence polarization reagent based on the peptide sequence 20–33 of ubiquitin. The peptide is modified on the *N*-terminus with a 5-carboxytetramethylrhodamine (TAMRA) dye and conjugated at Lys27 to Ub via a native isopeptide bond.

sequence

TAMRA-VKAKIQDK(Ub)EGIPPD

important: sample preparation

- dilute DMSO stock with milliQ and mix well (e.g., 24x to a 100 µM stock; this can be aliquoted and stored)
- For assays this 100 µM stock can be diluted for example 1000x in buffer affording a final assay solution of 100 nM. The DMSO concentration during the assay is now 0.01 vol%.
- the concentration can be verified by comparing the fluorescence intensity with that of a known concentration of TAMRA.
- for full experimental details please see reference 5.

Literature. (1) Tirat et al. *Analytical Biochem.* **2005**, *343*, 244. (2) Huang et al. *Methods in Mol Biol* **2009**, *565*, 127. (3) Levine et al. *Analytical Biochem.* **1997**, *247*, 83. (4) Faesen et al. *Chem. Biol.* **2011**, *18*, 1550. (5) Geurink et al. *ChemBiochem* **2012**, *13*, 293.

General experimental procedures for fluorescence polarization assays

FP assays were performed on a PerkinElmer Wallac EnVision 2010 Multilabel Reader with a 531 nm excitation filter and two 579 nm emission filters. Fluorescence intensities were measured in the S (parallel) and P (perpendicular) direction. FP values are given in mP (millipolarization) and calculated using the following formula (1):

$$\text{Polarization (mP)} = \frac{S-(G \cdot P)}{S+(G \cdot P)} \cdot 1000 \quad (1)$$

The confocal optics were adjusted with the average P and S values for TAMRA-Lys-Gly and the grating factor (G) was determined using a polarization value (L) for TAMRA of 50 mP using the following formula (2):

$$G = \frac{\text{average S}}{\text{average P}} \cdot \frac{1 - \left(\frac{L}{1000}\right)}{1 + \left(\frac{L}{1000}\right)} \quad (2)$$

The assays were performed in “non-binding surface flat bottom low flange” black 384-well plates (Corning) at room temperature in a buffer containing 20 mM Tris-HCl, pH 7.5, 5 mM DTT, 100 mM NaCl, 1 mg/mL 3-[(3-cholamidopropyl) dimethylammonio] propanesulfonic acid (CHAPS) and 0.5 mg/mL bovine gamma globulin (BGG). Each well had a volume of 20 μ L. Buffer and enzyme were predispensed and the reaction was started by the addition of substrate. Kinetic data was collected in intervals of 2.5 or 3 min. From the obtained polarization values (P_t) the amount of processed substrate (S_t) was calculated with to the following formula (3):

$$S_t = S_0 - S_0 \times \left[\frac{P_t - P_{\min}}{P_{\max} - P_{\min}} \right] \quad (3)$$

Where P_t is the polarization measured (in mP); P_{\max} is the polarization of 100% unprocessed substrate (determined for every reagent at all used substrate concentrations); P_{\min} is the polarization of 100% processed substrate; S_0 is the amount of substrate added to the reaction.

From the obtained P_t values the values for initial velocities (v_i) were calculated. The v_i values are used to determine the Michaelis-Menten constants (K_m , V_{\max} and k_{cat}) by fitting the data according to formula (4) (where $k_{\text{cat}} = V_{\max}/[E]$). All experimental data was processed using Ms Excel and Prism 4.03 (GraphPad Software, Inc.).

$$v_i = \frac{V_{\max} \times S_0}{K_m + S_0} \quad (4)$$