

UbiQ

targeting the ubiquitin system

Ub photoLeu8 (human sequence, Met1Nle, synthetic)

UbiQ code : UbiQ-153

Batch # : B01062019-001

Amount : 100 ug, lyophilized powder

Purity : $\geq 95\%$

Mol. Weight : 8.58 kDa

Storage : upon arrival, powder at -20°C ; solution at -80°C . Please avoid multiple freeze/thaw cycles and store dark.


Productsheet

Background. UbiQ-153 is a crosslinking reagent based on ubiquitin in which Leu8 is replaced by the photoreactive analog photoleucine (photoLeu, ^{PL}, Figure 1B). Upon irradiation of the diazirine with UV light, a very reactive carbene species is formed (Figure 1C). Unspecific binding is minimal for carbene based crosslinking because carbenes are quenched quickly by water, meaning only residues that are very nearby (and thereby contribute to tight binding) will react covalently.

A

NleQIFVKT^{PL}LGKTTITLEVEPSDTIENVKAKIQDKEGIPPDQQRLLIFAGKQLEDGRTLSDYNIQKESTLHLVLRRLGG

B



C




Figure 1. A: sequence; to avoid oxidation, Met1 has been replaced by its isostere norleucine (Nle). B: photoleucine. C: carbene formation.

important: sample preparation

- dissolve the powder in as little DMSO as possible (e.g. 20 - 40 mg/mL)
- add the DMSO stock slowly to milliQ (please note the order of addition)
- buffer the aq. solution as desired
- under normal laboratory light, UbiQ-153 is photo-stable
- for detailed experimental procedures please see reference 1.

Literature. (1) Chojnacki et al. *Cell Chem Biol* **2017**, *24*, 443. (2) Liang et al. *Angew Chem Int Ed* **2017**, *56*, 2744. (3) Zhou et al. *Nat Comm* **2016**, *7*, article 10589. (4) Dubinsky et al. *Bioorg Med Chem* **2012**, *20*, 554.

UbiQ Bio BV

Chamber of Commerce 50023438
VAT NL822502136B01

Science Park 301
1098 XH Amsterdam
The Netherlands

t +31 20 303 1970
e info@ubiqbio.com
i www.ubiqbio.com

Rabobank
IBAN: NL86 RABO 0150658907
BIC/SWIFT: RABONL2U