

**Ub K6C** (human sequence, recombinant)

UbiQ code : UbiQ-174 Batch # : B01092016-001

Amount : 1.00 mg, lyophilized powder

Purity : ≥95% by RP-HPLC

Mol. Weight: 8.54 kDa

Storage : powder at -20°C; solution at -80°C. Please avoid multiple freeze/thaw cycles.


## **Productsheet**

**Background.** UbiQ-174 is based on the ubiquitin protein (Ub) in which lysine 6 has been mutated to a cysteine residue. Ub Lys-to-Cys mutants can be used to prepare well-defined Ub-chains<sup>1,2</sup> and site-selective functionalized Ub proteins (using thiol reactive moieties).

## Α

MQIFVCTLTGKTITLEVEPSDTIENVKAKIQDKEGIPPDQQRLIFAGKQLEDGRTLSDYNIQKESTLHLVLRLRGG





**Figure 1.** A: sequence. B: LC-MS analysis. Mobile phase A = 1% CH<sub>3</sub>CN, 0.1% formic acid in water and B = 1% water and 0.1% formic acid in CH<sub>3</sub>CN. XBridge BEH300 C18 5 $\mu$ m 4.6x100mm; flow rate = 0.8 mL/min, runtime = 10 min, column T = 40°C. Gradient: 30-60% B over 6.5 min.

## important: sample preparation

- dissolve the powder in as little DMSO as possible (e.g. 20 40 mg/mL)
- add the DMSO stock slowly to milliQ (please note the order of addition)
- buffer the aq. solution as desired (e.g. 50 mM HEPES pH 8, 100 mM NaCl)
- for example, a final buffered stock of 0.5 mg/mL (59 uM) will contain 1.25 vol% DMSO when prepared from a 40 mg/mL DMSO stock.

Literature. (1) Raasi et al Methods Mol Biol. 2005, 301, 47. (2) Valkevich et al. J Am Chem Soc. 2012, 134, 6916.